

19



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

11 Publication number:

0 032 063  
A2

12

# EUROPEAN PATENT APPLICATION

21 Application number: 80304749.7

22 Date of filing: 30.12.80

51 Int. Cl.<sup>3</sup>: A 61 K 31/225

C 07 C 69/736, C 07 C 69/738  
C 07 C 149/40, C 07 C 79/46  
C 07 C 59/68, C 07 C 62/34  
C 07 C 147/11, C 07 C 43/205  
C 07 C 67/313, C 07 C 67/333

30 Priority: 02.01.80 US 108978

43 Date of publication of application:  
15.07.81 Bulletin 81/28

84 Designated Contracting States:  
AT BE CH DE FR GB IT LI LU NL SE



71 Applicant: WARNER-LAMBERT COMPANY  
201 Tabor Road  
Morris Plains, New Jersey 07950(US)

72 Inventor: Sircar, Ila  
3615 Charter Place  
Ann Arbor Michigan 48105(US)

74 Representative: Jones, Michael Raymond et al,  
Haseltine Lake & Co. 28 Southampton Buildings  
Chancery Lane  
London WC2A 1AT(GB)

54 Novel substituted arylene compounds and processes for their production.




57 Novel substituted arylene compounds and processes for their production are disclosed. These new compounds are useful as anti-arteriosclerotic agents. The novel compounds have the following formulae:-



EP 0 032 063 A2

Croydon Printing Company Ltd.

10/743,951 B04



The radicals X, Y, L and M are each oxygen, sulfur, sulfone or sulfoxide and the other substituents are as defined at length in claim 1.

-1-


NOVEL SUBSTITUTED ARYLENE COMPOUNDS AND PROCESSES FOR  
THEIR PRODUCTION

This invention relates to novel substituted  
arylene compounds and processes for their production.


Elevated levels of blood cholesterol and blood  
lipids are conditions which are believed related to the  
onset of arteriosclerosis. Thus, compounds capable of  
reducing the levels of these blood constituents are  
recognized as potentially useful anti-arteriosclerotic  
agents. The prior art contains many materials which  
are characterised as potentially useful anti-arteriosclerotic  
agents. These prior art agents and intermediates thereto,  
which the applicants believe are most closely structurally  
related to the compounds of the present invention are as  
follows:


United States Patent Specification No. 3,686,271  
discloses a class of arylene bis alpha-alkyloxyacetic acids  
and certain derivatives; and United States Patent  
Specification No. 3,769,436 discloses a class of  
arylenebisoxoacetic acids and certain derivatives.

One aspect of the present invention provides a  
compound having one of the formulae 1 to 5:-



-2-




wherein X and Y are the same or different and are oxygen, sulfur, sulfone or sulfoxide; m and m' are the same or different and are integers in the range from 2 to 5; each R is CH<sub>2</sub>OH, CHO or CO<sub>2</sub>A wherein A is a hydrogen atom, an alkyl radical having from 1 to 6 carbon atoms, an alkali or alkaline earth metal cation, an organic amine cation or an ammonium cation; each of R<sup>1</sup> and R<sup>2</sup>, which are the same or different, is an alkyl radical having from 1 to 6 carbon atoms, or R<sup>1</sup> when taken together with the R<sup>2</sup> which shares the same carbon atom represents -(CH<sub>2</sub>)<sub>3-5</sub>-; and R<sup>3</sup> is a hydrogen or halogen atom, an alkyl radical having from 1 to 6 carbon atoms, methoxy, phenyl, trifluoromethyl, 2,5-dichloro, 2,5-dimethyl, 2,3,5-trimethyl, carboxylic acyl or carboxylato alkyl of from 1 to 6 carbon atoms, alphahydroxyalkyl having from 1 to 6 carbon atoms, 2-chlorophenyl, 4-chlorophenyl, 2,4-dichlorophenyl, 4-methoxyphenyl, 2-tolyl, 4-tolyl or -D wherein D is O, CO, S, SO, SO<sub>2</sub>, CH<sub>2</sub>, or CHOH; provided that R<sup>3</sup> is not a hydrogen atom when all of the following conditions are satisfied, namely when X and Y are both oxygen atoms, m and m' are both 2, R<sup>1</sup> and R<sup>2</sup> are both CH<sub>3</sub>, and R is CO<sub>2</sub>H:



wherein L and M are the same or different and are oxygen, sulfur, sulfone or sulfoxide; n and n' are the same or different and are integers in the range from 2 to 5; each of R<sup>4</sup> and R<sup>5</sup> which are the same or different is an alkyl radical having from 1 to 6 carbon atoms; R' is CH<sub>2</sub>OH, CHO or CO<sub>2</sub>A wherein A is a hydrogen atom, an alkyl of from

-3-

1 to 6 carbon atoms, an alkali or alkaline earth metal cation, an organic amine cation or an ammonium cation; and  $R^6$  is a hydrogen atom, 4-formyl, 2-nitro, 2-methyl, 5-methyl, 4-benzoyl or 4-chloro:



- 5 wherein, in formula 3, 4 or 5, each  $R''$  is  $\text{CH}_2\text{OH}$ ,  $\text{CHO}$  or  $\text{CO}_2\text{A}$  wherein A is a hydrogen atom, an alkyl radical having from 1 to 6 carbon atoms, an alkali or alkaline earth metal cation, an organic amine cation or an ammonium cation;
- or, when  $R, R'$  or  $R''$  is  $\text{COOH}$  in any such compound,
- 10 a pharmaceutically acceptable salt of that compound.

The compounds of the present invention are useful as hypocholesterolemic agents and possess the additional advantage of elevating the high density lipoprotein fraction of cholesterol (HDL-cholesterol), which is known

15 to lower the risk factor of coronary heart disease


-4-




(Gordon, T. et al., High Density Lipoprotein as a Protective Factor Against Coronary Heart Disease, May 1977, The American Journal of Medicine, Vol. 62, pp. 707-714). Certain compounds of the invention also reduce the  
5 low density lipoprotein fraction of cholesterol (LDL-cholesterol), thus further reducing the risk factor of coronary heart disease.


The invention also relates to methods for using the above described compounds as anti-arteriosclerotic  
10 agents for treating mammals; additionally the present invention also provides pharmaceutical compositions comprising a plasma high density lipoprotein increasing amount of a compound as described above as well as the pharmaceutically acceptable acid-addition salts thereof  
15 and mixtures thereof.

The compounds of the present invention may be prepared by several processes in accordance with another aspect of the present invention. These processes, which are equivalent for the purposes of the  
20 present invention, are described hereinbelow with specific reference to the dioxyphenylene compounds of the invention. This description is utilized merely for convenience and those skilled in the art will recognize that the analogous sulfur-containing compounds can be  
25 prepared by substituting sulfur-containing starting materials or utilizing substantially identical reactions or obvious variations thereof. Those skilled in the art will also recognize that the sulfur compounds can be utilized to prepare corresponding sulfone and sulfoxide  
30 compounds by art-recognized oxidation techniques.


In the preferred method of the invention, a compound of the formula I



wherein (Ar) is , , or ; and R<sub>x</sub> represents any of the hereinbefore defined substituents R<sup>3</sup> and R<sup>6</sup>, is reacted with a compound (an alkylating agent) of the formula II




II

wherein n, R, R<sup>1</sup> and R<sup>2</sup> are as hereinabove defined. In formula I (Ar) is preferably , and in formula II n is preferably 3, R is preferably CO<sub>2</sub>CH<sub>3</sub> and R<sup>1</sup> and R<sup>2</sup> are preferably CH<sub>3</sub>.

Q is a substituent known to those skilled in the art as a "leaving group" [C.B. Snyder and A.R. Soto, J. Org. Chem., 30, 673 (1965)]. For purposes of the present invention, the halogens are suitable leaving groups and the preferred halogen is bromine. Other suitable leaving groups such as p-toluenesulfonyloxy will be known to those skilled in the art. The reaction is usually carried out in the presence of a base, examples of which are the alkali metal hydrides, amides, hydroxides, alkoxides and carbonates such as sodium hydride, potassium amide, potassium hydroxide, sodium methoxide, potassium t-butoxide and sodium carbonate. The preferred bases are the alkali metal hydrides, especially sodium hydride. The reaction is conducted usually in an anhydrous, non-protic solvent such as an ether, a hydrocarbon or a tertiary amide; examples of such solvents are dimethoxyethane, dioxane, tetrahydrofuran, pentane, cyclohexane, benzene, xylene, and N,N-dimethylformamide. Other suitable solvents will suggest themselves to those skilled in the art. The preferred solvent is N,N-dimethylformamide. The time and temperature ranges for conducting the



foregoing reaction may be varied, and in general, a temperature range of about 20°C to about 70°C and a time range of about 2 to about 24 hours can be used. The preferred ranges are about 60°-70°C for about 4-5 hours. The ranges can vary depending on the reactants, the solvent and the base being utilized, and it is within the skill of the art to choose these ranges to optimize the yield for a specific reaction. When carrying out the above-described reaction, it is preferred to use slightly more than two equivalents of the alkylating agent, II, and slightly more than two equivalents of base per mole of the compound I. The use of a greater excess of base, for example from 4 to 10 equivalents, is desirable when an alkali metal carbonate such as sodium or potassium carbonate is utilized. The product of the reaction, represented by the preferred structure III,




may be isolated by standard procedures, for example, addition of the reaction mixture to water, separating the organic phase when present, extracting the aqueous phase with a water-immiscible solvent such as diethyl ether, washing the combined organic phases with water, drying the washed organic phase with, for example,  $\text{MgSO}_4$ , evaporating the solvent and purifying the product by, for example, vacuum distillation. Alternatively, when R is an ester this function may be saponified, with, for example, methanolic sodium

hydroxide, and the acid product may be isolated and purified by well-known methods. If desired, the ester functions may be saponified after the ester has been isolated and purified. The above described reaction of compounds I and II will produce products, III, wherein  $n$  and  $n'$  are the same. When it is desired to prepare a compound of formula III wherein  $n$  and  $n'$  are different, those skilled in the art will understand that the reaction can be performed in two separate stages, i.e., each step utilizing substantially one equivalent of alkylating agent II, and substantially one equivalent of base for each mole of compound I.


The compounds I, and the alkylating agents II, are either commercially available or are readily prepared. The preferred alkylating agent, methyl 5-bromo-2,2-dimethylpentanoate may be prepared from 2,2-dimethyl-4-pentenoic acid [J. Am. Chem. Soc., 77, 1092 (1955)] by the addition of hydrogen bromide in the presence of benzoyl peroxide.

A second process for preparing the compounds of the invention comprises the reaction of an arylenebis(oxy)alkyl compound of formula IV




IV

wherein  $\text{Ar}$ ,  $n$ ,  $n'$ ,  $\text{R}_x$  and  $\text{Q}$  are as previously defined with a compound of formula V



V


wherein R, R<sup>1</sup> and R<sup>2</sup> are as previously defined. Z is an alkali metal which is preferably lithium, R is preferably CO<sub>2</sub>A', wherein A' is sodium or lithium, and (Ar) is preferably .

Preferred solvents for use in this reaction are anhydrous polar solvents such as tetrahydrofuran, tetrahydropyran, dimethoxy ethane, diethylene glycol dimethyl ether and dimethyl sulfoxide. Tetrahydrofuran is the preferred solvent at temperatures, up to about 40°C whereas diethylene glycol dimethyl ether is the preferred solvent at temperatures above 40°C. The reaction solvent can also be a hydrocarbon such as pentane, heptane, benzene or toluene, as well as mixtures thereof.

The time and temperature for conducting the reaction are dependent on the particular reactants and the solvent that is used. In general a temperature from about -10°C to about 60°C and a reaction time of from about one hour to about 24 hours can be used. According to the preferred method, the reaction is carried out at a temperature between 0°C to about 30°C for about 16 hours. When reacting compounds IV and V, a ratio of 2 equivalents of V to 1 mole of IV is


normally utilized, but it is preferred to use a slight excess of the arylenebis(oxy)alkyl compound, IV.

The product may be isolated and purified by standard procedures, such as those described hereinabove for compound III. When the product is in the form of the diester, it may be saponified, if desired, for example with dilute base either before or after its isolation and purification. The starting arylenebis(oxy)alkyl compounds, IV, may be prepared by well-known methods such as by reaction of the corresponding compound, I, with a compound of the formula VI [R. Adams and L.N. Whitehill, J. Am. Chem. Soc., 63, 2073 (1949)],



# VI

wherein Q and Q' are leaving groups which may be the same or different and are defined hereinabove, n is also as previously defined. For purposes of the invention, Q and Q' are preferably different, for example Q may be chlorine when Q' is bromine. The reaction of compound I with VI is performed in substantially the same manner as that described hereinabove for the reaction of compound II with compound I.

Certain of the compounds may be prepared by aromatic ring substitution. Thus, the benzene ring of a compound of formula III wherein  $\text{Ar}$  is  and R<sub>x</sub> is hydrogen may be, for example, halogenated.

The compounds of the invention wherein R is CHO may be prepared by reduction of the corresponding acid or by oxidation of the corresponding alcohol. These compounds can also be prepared by using starting materials such as II or V wherein R is a protected CHO group such as in the form of an ethylene acetal. The protecting group is subsequently removed, for example, by dilute acid hydrolysis of the ethylene acetal.

The compounds of the invention wherein R is  $\text{CH}_2\text{OH}$  can also be prepared from the corresponding aldehyde, free acid or ester by, for example, reduction with lithium aluminum hydride. The carboxylic acids of the invention, i.e., the compounds of formula III wherein R, R' and R'' are  $\text{CO}_2\text{H}$ , form carboxylate salts with any of a variety of inorganic and organic bases. Suitable bases for purposes of the invention, are those which form pharmaceutically-acceptable salts such as sodium hydroxide, sodium carbonate, sodium bicarbonate, potassium hydroxide, calcium hydroxide, ammonia and amines. The salts are converted back to their respective carboxylic acids by treatment with an acid such as dilute hydrochloric acid. The carboxylic acids and their respective salts differ in certain physical properties such as solubility properties but they are otherwise equivalent for purposes of the invention.

The compounds of the invention can exist in unsolvated as well as solvated forms, including hydrated forms. In general, these solvated forms with pharmaceutically acceptable solvents such as water, ethanol and the like are equivalent to the unsolvated forms for purposes of the invention.

The compounds of the invention are new chemical substances, of value as pharmacological agents which increase the high density lipoprotein fraction of cholesterol (HDL-cholesterol) at the expense of the low density lipoprotein fraction of cholesterol

(LDL-cholesterol). In addition, many of the compounds of the invention reduce total plasma cholesterol levels. The risk of coronary heart disease is thereby reduced. The effectiveness of the compounds of the invention was established by the screening procedure described in Maxwell, R.E., Nawrocki, J.W., and Uhlenendorf, P.D., *Artery*, 4, 303 (1978).

A compound is considered active in this procedure if it increases the HDL fraction of cholesterol by at least 50%. The test results obtained for compounds of the invention are shown in the following table.

#### TEST RESULTS

Compound of Example	Dose mg/kg	Percent Change		
		Plasma Cholesterol	HDL Cholesterol	HDL/LDL Electro-Phoretic
1A	50	-30	347	435
1B	50	0	381	318
1D	50	-30	385	595
1G	50	0	480	-
1K	50	0	628	-
1M	50	0	674	-
1O	50	0	723	-
1P	50	+39	1100	-
1Q	50	0	421	-
1R	50	-30	686	-
1S	50	+32	428	-
2	50	-26	566	871
2A	50	-37	743	1009
2B	50	-45	728	1311
4	50	-27	1410	2432
4	25	-	741	1798
4	12.5	-	518	945

TEST RESULTS (Continued)

Compound of Example	Dose mg/kg	Percent Change			HDL/LDL Electro- Phoretic
		Plasma Cholesterol	HDL Cholesterol		
5	50	-45	791		2472
5A	50	0	1477		1722
5D	50	-55	384		1091
5E	50	-62	348		377
5F	50	-53	753		1443
5F	25	-	206		493
5F	12.5	-	94		242
5I	50	-52	294		794
5J	50	-28	383		510
5K	50	-37	317		556
5L	50	0	276		274
5M	50	-44	628		785
5M	25	-	232		302
5M	12.5	-	74		101
6	50	-33	289		388
7	50	0	907		703
8	50	-27	525		509
10	50	-44	389		-
11	50	-	410		-
12	50	+40	319		-
13	50	+61	1292		-
14	-	-	-		-
17	50	0	947		-

The compounds of the invention can be administered either orally or parenterally. They can be combined with a solid or liquid carrier or diluent and made available in varying amounts in such pharmaceutical forms as tablets, capsules, powders, and aqueous and non-aqueous suspensions and solutions. The preferred daily dosage is from about 10 to about 250 mg/kg which may be administered in either single or divided dosage units.

The starting materials for preparing the compounds of the invention are either commercially available or can be readily prepared by published literature procedures, for example:

2,5-Dioxybenzophenone is prepared according to J. Herzig and Br. Hofmann, Ber, 41, 143 (1908);

Phenylthio-benzohydroquinone is prepared according to O. Dimroth, L. Kraft, and K. Aichinger, Ann. der Chemie, 545, 124 (1940). All the substituted phenylhydroquinones are prepared according to D.E. Kvalnes, J. Am. Chem. Soc., 56, 2478 (1934). Fluorohydroquinone and trifluoromethylhydroquinone are prepared according to the procedure of A.E. Feiring, and W.A. Sheppard, J. Org. Chem., 40, 2543 (1975).

The invention is illustrated by the following examples.

#### EXAMPLE 1

A solution of 3.32 g of *t*-butylhydroquinone in 15 ml of dimethylformamide is added dropwise to a suspension of 2.4 g of 50% sodium hydride in 25 ml of dimethylformamide, and allowed to stir for 1/2 an hour at room temperature. 11.2 g of methyl 5-bromo-2,2-dimethylpentanoate is added followed by heating the mixture at 60-70°C for 6 hrs. Dimethylformamide is distilled off, the residue is poured into water and the organic material is extracted with ether. The ether extract is washed with water until neutral, dried over anhydrous magnesium sulfate and the ether is evaporated. The residue is purified by distillation



yielding the product, dimethyl 5,5'-[[2-(1,1-dimethylethyl)1,4-phenylene]bis(oxy)]bis[2,2-dimethylpentanoate] b.p. 200°C at 0.5 mm of Hg; yield 4g. .

Following the general procedure of Example 1, with the substitution of an equivalent amount of the specified diphenol, the following products are obtained:

A. From fluorohydroquinone, the product obtained is dimethyl 5,5'-[(2-fluoro-1,4-phenylene)bis(oxy)]bis[2,2-dimethylpentanoate], m.p. 33-35° following crystallization from ether-petroleum ether.

B. From 2,5-dihydroxyacetophenone, the product obtained is dimethyl 5,5'-[(2-acetyl-1,4-phenylene)-bis(oxy)]bis[2,2-dimethylpentanoate] b.p. 210°C at 0.1 mm of Hg.

C. From 2,4-dihydroxybenzophenone, the product obtained is dimethyl 5,5'-[(4-benzoyl-1,3-phenylene)bis(oxy)]bis[2,2-dimethylpentanoate], which is hydrolysed directly by refluxing with a solution of 2N methanolic sodium hydroxide to give the corresponding acid, 5,5'-[(4-benzoyl-1,3-phenylene)bis(oxy)]bis[2,2-dimethylpentanoic acid] m.p. 125-126°C, following crystallization from tetrahydrofuran-isopropyl ether.

D. From 2-methylresorcinol, the product obtained is dimethyl 5,5'-[(2-methyl-1,3-phenylene)bis(oxy)]bis[2,2-dimethylpentanoate], b.p. 200°C at 0.3 mm of Hg.

E. From 5-methylresorcinol, the product obtained is dimethyl 5,5'-[(5-methyl-1,3-phenylene)bis[2,2-dimethylpentanoate], b.p. 200°C at 0.3 mm of Hg.

F. From 4-chlorobenzene 1,3-dithiol the product obtained is dimethyl 5,5'-[(4-chloro-1,3-phenylene)-bis(thio)]bis[2,2-dimethylpentanoate] b.p. 200°C at 0.02 mm of Hg.

G. From o-chlorophenylhydroquinone, the product obtained is 5,5'-[(2'-chloro[1,1'-biphenyl]-2,5-diyl)-bis(oxy)]bis[2,2-dimethylpentanoic acid], m.p. 101-2°C, following crystallization from isopropyl ether.

- H. From *p*-anisylhydroquinone, the product obtained is 5,5'-[(4'-methoxy[1,1'-biphenyl]-2,5-diyl)bis(oxy)]-bis[2,2-dimethylpentanoic acid], m.p. 69-73° (d), following crystallization from isopropyl ether.
- I. From *p*-chlorophenylhydroquinone, the product obtained is 5,5'-[(4'-chloro[1,1'-biphenyl]-2,5-diyl)-bis(oxy)]bis[2,2-dimethylpentanoic acid], m.p. 110-111°C, following crystallization from tetrahydrofuran-petroleum ether.
- J. From *p*-tolylhydroquinone, the product obtained is 5,5'-[(4'-methyl[1,1'-biphenyl]-2,5-diyl)bis(oxy)]-bis[2,2-dimethylpentanoic acid], m.p. 65-8°C, following crystallization from chloroform.
- K. From *o*-tolylhydroquinone, the product obtained is 5,5'-[(2'-methyl[1,1'-biphenyl]2,5-diyl)bis(oxy)]-bis[2,2-dimethylpentanoic acid], m.p. 65-8°C, following crystallization from isopropyl ether.
- L. From 2,5-dihydroxybenzaldehyde, the product obtained is dimethyl 5,5'-[(2-formyl-1,4-phenylene)-bis(oxy)]bis[2,2-dimethylpentanoate], b.p. 220°C at 0.1 mm of Hg.
- M. From trifluoromethylhydroquinone, the product obtained is 5,5'-[(2-trifluoromethyl-1,4-phenylene)bis(oxy)]bis[2,2-dimethylpentanoic acid], m.p. 114-5°C, following crystallization from tetrahydrofuran-hexane.
- N. From 2,4-dichlorophenylhydroquinone, the product obtained is 5,5'-[(2',4'-dichloro[1,1'-biphenyl]2,5-diyl)-bis(oxy)]bis[2,2-dimethylpentanoic acid], m.p. 82-6°C, following crystallization from isopropyl ether.
- O. From 2,5-dihydroxypropiophenone, the product obtained is dimethyl 5,5'-[[2-(1-oxopropyl)-1,4-phenylene]bis(oxy)]bis[2,2-dimethylpentanoate].
- P. From 2,5-dihydroxybutyrophenone, the product obtained is dimethyl 5,5'-[[2-(1-oxobutyl)-1,4-phenylene]bis(oxy)]bis[2,2-dimethylpentanoate].

Q. From 2,5-dihydroxybenzophenone, the product obtained is dimethyl 5,5'-[(2-benzoyl-1,4-phenylene)-bis(oxy)]bis[2,2-dimethylpentanoate].

R. From 2,5-dimethylhydroquinone, the product obtained is dimethyl 5,5'[2,5-dimethyl-1,4-phenylenebis(oxy)]-bis[2,2-dimethylpentanoate], m.p. 60-2°C, following crystallization from isopropyl ether.

S. From phenylthiobenzohydroquinone, the product obtained is 5,5'-[(2-phenylthio)-1,4-phenylene]bis-(oxy)]bis[2,2-dimethylpentanoic acid], m.p. 117-8°C, following crystallization from tetrahydrofuran-isopropyl ether.

#### EXAMPLE 2

A solution of 3.5 g of 2-methoxyhydroquinone in 35 ml of methanol is stirred with 3.0 g of sodium methoxide for thirty minutes. 12.3 g of methyl 5-bromo-2,2-dimethylpentanoate is added and the mixture is refluxed for 24 hours. Methanol is removed under reduced pressure and the residue is poured into water. An oil separates, which is extracted with ether and the ether extract is washed with water, dried over anhydrous magnesium sulfate and the ether is evaporated. The residue is purified by distillation yielding the product, dimethyl 5,5'-[(2-methoxy-1,4-phenylene)bis(oxy)]bis[2,2-dimethylpentanoate] b.p. 200°C at 0.2 mm of Hg; yield 6.6 g.

Following the general procedure of Example 2, with the substitution of an equivalent amount of the specified diphenol, the following products are obtained:

A. From methylhydroquinone the product obtained is dimethyl 5,5'-[(2-methyl-1,4-phenylene)bis(oxy)]-bis[2,2-dimethylpentanoate] b.p. 192°C at 0.15 mm of Hg.

B. From bromohydroquinone, the product obtained is dimethyl 5,5'[(2-bromo-1,4-phenylene)bis(oxy)]bis[2,2-dimethylpentanoate] b.p. 200°C at 0.3 mm of Hg.

EXAMPLE 3

A suspension of 3.45 g of 2,4-dihydroxybenzaldehyde, 10.4 g of anhydrous potassium carbonate, 0.5 g of potassium iodide and 13.4 g of methyl 5-bromo-2,2-dimethylpentanoate in 70 ml of acetone is refluxed for 24 hours. The suspension is cooled, filtered and the solvent is evaporated. The residue is poured into water, and the organic material is extracted with ether. The ether extract is washed with water, dried over anhydrous magnesium sulfate and evaporated. The residue is purified by distillation yielding the product, dimethyl 5,5'-[(4-formyl-1,3-phenylene)bis(oxy)]bis[2,2-dimethylpentanoate], b.p. 210°C at 0.1 mm of Hg; yield 6.5g.

A. From 2-nitroresorcinol the product obtained is dimethyl 5,5'-[(2-nitro-1,4-phenylene)bis(oxy)]bis[2,2-dimethylpentanoate] b.p. 230°C at 1.0 mm of Hg.

B. From 4,4'-biphenyldithiol, the product obtained is dimethyl 5,5'-[[1,1'-biphenyl]-4,4'-diylbis(thio)]bis[2,2-dimethylpentanoate], m.p. 46-7°, following crystallization from isopropyl ether.

EXAMPLE 4

A solution of 60.0 g of phenylhydroquinone in 200 ml of dimethylformamide is added dropwise to a suspension of 27.5 g of 59% sodium hydride in 100 ml of dimethylformamide with stirring under N<sub>2</sub> blanket. The mixture is stirred for one hour at room temperature to complete the reaction. One hundred fifty-six ml of methyl 5-bromo-2,2-dimethylpentanoate is added dropwise at room temperature followed by heating the mixture at 80-85°C for 12-14 hours. The mixture is cooled, filtered from the inorganic residue and dimethylformamide is distilled under reduced pressure. The residue is treated with water and the organic material is extracted with ether. The ether extract is dried over anhydrous magnesium sulfate and ether removed leaving behind an oil. The unreacted alkyl bromide is removed

by distillation under high vacuum and the residue is purified by chromatography over silica gel. The crude diester (160 g) is directly hydrolyzed by refluxing with 1600 ml of 2N methanolic sodium hydroxide for 3 hours. Methanol is removed and the residue is treated with water followed by acidification. The product obtained is 5,5'-[(1,1'-biphenyl)-2,5-diylbis(oxy)]-bis[2,2-dimethylpentanoic acid], mp 139-140°C, following crystallization from tetrahydrofuran-isopropyl ether, yield 125 g.

#### EXAMPLE 5

17.6 g of isobutyric acid is added dropwise to a stirred suspension of 10.6 g of 50% NaH and 20.6 g of diisopropylamine in 170 ml of anhydrous tetrahydrofuran at room temperature under nitrogen. The mixture is refluxed for 30 minutes followed by cooling to 0°C. At this time 138 ml of a solution of butyllithium in heptane is added slowly. The ice-bath is retained for 30 minutes followed by warming to 30-40° for another 30 minutes. The slightly turbid solution is cooled to 0°C and a solution of 26.3 g of 1,4-bis(3-chloropropoxy)-benzene in 50 ml of dry tetrahydrofuran is added dropwise while the temperature was maintained below 10°C. After 30 minutes the mixture is allowed to warm to room temperature and stirring is continued for 16 hours. The mixture is cooled and hydrolysed with 250 ml of water and the aqueous phase is separated, washed with 100 ml of ether, and acidified with 6N hydrochloric acid yielding the product, 5,5'-[1,4-phenylenebis(oxy)]bis[2,2-dimethyl pentanoic acid], m.p. 168°C, following crystallization from tetrahydrofuran-isopropyl ether, yield 23 g.

Following the general procedure of Example 5, with the substitution of an equivalent amount of the specified starting material the following products are obtained:

- A. From 1,4 bis(4-chlorobutoxy)benzene the product obtained is 6,6'-[1,4-phenylenebis(oxy)]bis[2,2-dimethylhexanoic acid], m.p. 158-159°C following crystallization from tetrahydrofuran-hexane.
- B. From cyclobutane carboxylic acid the product obtained is 1,1'-[1,4-phenylenebis[oxy(4,1-butanediyl)]]biscyclobutanecarboxylic acid, m.p. 132-133°C following crystallization from tetrahydrofuran-petroleum ether.
- C. From 1,4 bis(3-chloropropoxy)2,3,5-trimethylbenzene the product obtained is 5,5'-[(2,3,5-trimethyl-1,4-phenylene)bis(oxy)]bis[2,2-dimethylpentanoic acid], m.p. 143°C following crystallization from diethyl ether.
- D. From 1,4 bis(3-chloropropoxy)2,5 dichlorobenzene, the product obtained is 5,5'-[(2,5-dichloro-1,4-phenylene)bis(oxy)]bis[2,2-dimethylpentanoic acid], m.p. 162°C following crystallization from tetrahydrofuran-isopropyl ether.
- E. From 1,4 bis(2-chloroethoxy)-2-chlorobenzene, the product obtained is 4,4'-[(2-chloro-1,4-phenylene)bis(oxy)]bis[2,2-dimethylbutanoic acid], m.p. 135-137°C following crystallization from tetrahydrofuran - isopropyl ether.
- F. From 1,4 bis(3-chloropropoxy)-2-chlorobenzene, the product obtained is 5,5'-[(2-Chloro-1,4-phenylene)bis(oxy)]bis[2,2-dimethylpentanoic acid], m.p. 135°C following crystallization from tetrahydrofuran-isopropyl ether.
- G. From 4,4'-bis(3-chloropropoxy)1,1'biphenyl, the product obtained is 5,5'-[(1,1'-biphenyl)-4,4'-diylbis(oxy)]bis[2,2-dimethylpentanoic acid], m.p. 201°C, following crystallization from tetrahydrofuran-hexane.
- H. From 1,5-bis(3-chloropropoxy)naphthalene, the product obtained is 5,5'-[1,5-naphthalene-diylbis(oxy)]bis[2,2-dimethylpentanoic acid], m.p. 219°C following crystallization from dimethylformamide.

I. From 1,3 bis(3-chloropropoxy)benzene, the product obtained is 5,5'-[1,3-phenylenebis(oxy)]bis[2,2-dimethylpentanoic acid], m.p. 135-136°C, following crystallization from isopropyl ether.

J. From 1,3 bis(2-chloroethoxy)benzene, the product obtained is 4,4'-[1,3-phenylenebis(oxy)]bis[2,2-dimethylbutanoic acid], m.p. 104-105°C, following crystallization from diethyl ether-petroleum ether.

K. From 1,3 bis(4-chlorobutoxy)benzene, the product obtained is 6,6'-[1,3-phenylenebis(oxy)]bis[2,2-dimethylhexanoic acid], m.p. 85-86°C, following crystallization from isopropyl ether.

L. From 1,2 bis(5-chloropentoxy)benzene, the product obtained is 7,7'-[1,2-phenylenebis(oxy)]bis[2,2-dimethylheptanoic acid], m.p. 61°C, following crystallization from diethyl ether-petroleum ether.

M. From 1-(3-chloropropoxy)-4-[(3-chloropropyl)thio]benzene, the product obtained is 5-[(4-[(4-carboxy-4-methylpentyl)oxy]phenyl)thio]-2,2-dimethylpentanoic acid, m.p. 112°C following crystallization from chloroform-petroleum ether.

#### EXAMPLE 6

A solution of 3.05 g of 5-[[4-(4-carboxy-4-methylpentyl)oxy]-2,5-dimethylphenyl]thio]-2,2-dimethylpentanoic acid in 10 ml of glacial acetic acid containing 0.9 ml of hydrogen peroxide is stirred at room temperature for 6 hrs. The solution is poured into water and extracted with ether. The ether extract is washed with water, dried over anhydrous magnesium sulfate and the solvent is evaporated. The residual oil solidifies on trituration with isopropyl ether and the product obtained is 5-[[4-[(4-carboxy-4-methylpentyl)oxy]phenyl]sulfinyl]-2,2-dimethylpentanoic acid, m.p. 98-100°C, following crystallization from tetrahydrofuran-isopropyl ether; yield 1.9 g.

EXAMPLE 7

A solution of 6.56 g. of 6,6'-[1,4-phenylene bis(oxy)]bis[2,2-dimethylhexanoic acid] in 25 ml of tetrahydrofuran is added to a stirred suspension of 1.1 g of lithium aluminum hydride in 75 ml of tetrahydrofuran followed by refluxing for 5-6 hours. The mixture is cooled, and the excess hydride is decomposed with saturated sodium sulfate solution, filtered from the inorganic solids and the filtrate is evaporated. The product obtained is, 6,6'-[1,4-phenylenebis(oxy)]bis[2,2-dimethyl-1-hexanol], m.p. 72°C following crystallization from tetrahydrofuran-hexane, yield 4 g.

EXAMPLE 8

4.0 g of dimethyl 5,5'-[(1-acetyl-1,4-phenylene)-bis(oxy)]bis[2,2-dimethylpentanoate] is treated with 0.4 g of sodium borohydride in 40 ml of methanol at room temperature. After usual workup, the residue is hydrolysed with 2N methanolic sodium hydroxide yielding the product, 5,5'-[2-(hydroxyethyl)-1,4-phenylenebis(oxy)]bis[2,2-dimethylpentanoic acid], m.p. 141-143°C, following crystallization from tetrahydrofuranisopropyl ether, yield 1.2 g.

EXAMPLE 9

A solution of 2.5 g of dimethyl 5,5'-[[2-(1-hydroxypropyl)-1,4-phenylene]bis(oxy)]bis[2,2-dimethylpentanoate] in 25 ml of benzene was refluxed 8 hrs in the presence of catalytic amount of p-toluenesulfonic acid. After usual work up, the product obtained is dimethyl 5,5'-[2-(1-propenyl)-1,4-phenylenebis(oxy)]bis[2,2-dimethylpentanoate], yield 1.1 g.



EXAMPLE 10

3.0 g of dimethyl 5,5'-[(2-benzoyl-1,4-phenylene)-bis(oxy)]bis[2,2-dimethylpentanoate] is reduced catalytically and the crude ester saponified. The product thus obtained is 5,5'-[[2-(benzyl)-1,4-phenylene]bis(oxy)]bis[2,2-dimethylpentanoic acid], m.p. 140-141°C following crystallization from tetrahydrofuran-petroleum ether, yield 2.1 g.

EXAMPLE 11

A solution of 2.09 g of dimethyl 5,5'-[(2-benzoyl-1,4-phenylene)bis(oxy)]bis[2,2'-dimethylpentanoate] in 20 ml of 95% alcohol is refluxed with 2.0 g of Zn-dust and 2.0 g of sodium hydroxide. The suspension is filtered, acidified, and the product obtained is 5,5'-[2-(hydroxyphenylmethyl)-1,4-phenylenebis(oxy)]-bis[2,2-dimethylpentanoic acid], m.p. 125-128°C, following crystallization from ether-hexane, yield 1 g.

EXAMPLE 12

3.8 g of dimethyl 5,5'-[(2-formyl-1,4-phenylene)bis(oxy)]bis[2,2-dimethylpentanoate] is treated with 0.5 g of sodium borohydride in 40 ml of methanol at room temperature. After usual work up, the residue is hydrolysed with 2N methanolic sodium hydroxide and the product obtained is 2.7 g of 5,5'-[(2-hydroxymethyl)-1,4-phenylenebis(oxy)]bis[2,2-dimethylpentanoic acid], m.p. 128-9°C, following crystallization from tetrahydrofuran-isopropyl ether, yield 2.7 g.

EXAMPLE 13

3.65 g of dimethyl 5,5'-[[2-(1-oxobutyl)-1,4-phenylene]bis(oxy)]bis[2,2-dimethylpentanoate] is reduced with 0.5 g of sodium borohydride in 40 ml of methanol at room temperature. After usual work up, the residue is hydrolysed with 2N methanolic sodium hydroxide and the product obtained is 5,5'-[[2-(1-hydroxybutyl)-1,4-phenylene]bis(oxy)]bis[2,2-dimethylpentanoic acid], m.p. 122-123°C, following crystallization from ether-isopropyl ether, yield 2.1 g.

EXAMPLE 14

A solution of 2.8 g of 5,5'-[(2-phenylthio)-1,4-phenylene]bis(oxy)]bis[2,2-dimethylpentanoic acid] in 25 ml of methylene chloride is stirred overnight with 1.0 g of *m*-chloroperbenzoic acid at room temperature. The separated solid is filtered, washed with methylene chloride and dried. The product obtained is 5,5'-[[2-(phenylsulfinyl)-1,4-phenylene]bis(oxy)]bis[2,2-dimethylpentanoic acid], m.p. 141-3°C, following crystallization from tetrahydrofuran-isopropyl ether, yield 1.3 g.

EXAMPLE 15

2.4 g of 5,5'-[(2-(phenylthio)-1,4-phenylene]bis(oxy)]bis[2,2-dimethylpentanoic acid] is treated with 1.8 g of *m*-chloroperbenzoic acid in 75 ml of methylene chloride. The product obtained is 5,5'-[[2-(phenylsulfonyl)-1,4-phenylene]bis(oxy)]bis[2,2-dimethylpentanoic acid], m.p. 194-6°C, following crystallization from tetrahydrofuran-isopropyl ether, yield 1.8 g.

EXAMPLE 16

A solution of 4.77 g of 5-[[4-[(4-carboxy-4-methylpentyl)oxy]-2,5-dimethylphenyl]thio]-2,2-dimethylpentanoic acid in 20 ml of glacial acetic acid is treated with 4.9 ml of 30% hydrogen peroxide at 50°C in three portions followed by refluxing for two hours. The solution is poured into water and the solids are filtered, washed with water and dried. The product obtained is 5-[[4-[(4-carboxy-4-methylpentyl)oxy]phenyl]sulfonyl]-2,2-dimethylpentanoic acid, m.p. 106-7°C, following crystallization from tetrahydrofuran-isopropyl ether, yield 4.3 g.


EXAMPLE 17


8.2 g of dimethyl 5,5'-[[2-(1-oxopropyl)-1,4-phenylene]bis(oxy)]bis[2,2-dimethylpentanoate] is treated with 0.85 g of sodium-borohydride in 40 ml of methanol at room temperature. After usual work up, the


residue is hydrolysed with 2N methanolic sodium hydroxide and the product is 5,5'-[2-(1-hydroxypropyl)-1,4-phenylenebis(oxy)]bis[2,2-dimethylpentanoic acid], m.p. 112-113°C, following crystallization from tetrahydrofuran-petroleum ether, yield 4.1 g.

CLAIMS (for all States other than Austria)



1. A compound having one of the following formulae 1 to 5:-




wherein X and Y are same or different and are oxygen, sulfur, sulfone or sulfoxide; m and m' are the same or different and are integers in the range from 2 to 5; each R is CH<sub>2</sub>OH, CHO or CO<sub>2</sub>A wherein A is a hydrogen atom, an alkyl radical having from 1 to 6 carbon atoms, an alkali or alkaline earth metal cation, an organic amine cation or an ammonium cation; each of R<sup>1</sup> and R<sup>2</sup>, which are the same or different, is an alkyl radical having from 1 to 6 carbon atoms, or R<sup>1</sup> when taken together with the R<sup>2</sup> sharing the same carbon atom represents -(CH<sub>2</sub>)<sub>3-5</sub>-; and R<sup>3</sup> is a hydrogen or halogen atom, an alkyl radical having from 1 to 6 carbon atoms, methoxy, phenyl, trifluoromethyl, 2,5-dichloro, 2,5-dimethyl, 2,3,5-trimethyl, a carboxylic acyl or carboxylato alkyl of from 1 to 6 carbon atoms, alpha-hydroxyalkyl having from 1 to 6 carbon atoms, 2-chlorophenyl, 4-chlorophenyl, 2,4-dichlorophenyl, 4-methoxyphenyl, 2-tolyl, 4-tolyl or -D- wherein D is O, CO, S, SO, SO<sub>2</sub>, CH<sub>2</sub> or CHOH; provided that R<sup>3</sup> is not a hydrogen atom when all of the following conditions are satisfied, namely when X and Y are both oxygen atoms, m and m' are both 2, R<sup>1</sup> and R<sup>2</sup> are both CH<sub>3</sub>, and R is CO<sub>2</sub>H:



wherein L and M are the same or different and are oxygen, sulfur, sulfone or sulfoxide; n and n' are the same or different and are integers in the range from 2 to 5; each of R<sup>4</sup> and R<sup>5</sup> which are the same or different is an alkyl radical having from 1 to 6 carbon atoms; each R' is CH<sub>2</sub>OH, CHO or CO<sub>2</sub>A wherein A is a hydrogen atom, an alkyl radical having from 1 to 6 carbon atoms, an alkali or alkaline earth metal cation, an organic amine cation or an ammonium cation; and R<sup>6</sup> is a hydrogen atom, 4-formyl, 2-nitro, 2-methyl, 5-methyl, 4-benzoyl or 4-chloro:




-28-




wherein, in formula 3, 4 or 5, each  $\text{R}''$  is  $\text{CH}_2\text{OH}$ ,  $\text{CHO}$  or  $\text{CO}_2\text{A}$  wherein A is a hydrogen atom, an alkyl radical having from 1 to 6 carbon atoms, an alkali or alkaline earth metal cation, an organic amine cation or an ammonium cation:

or, when R, R' or R'' is  $\text{COOH}$  in any such compound, a pharmaceutically acceptable salt of that compound.


2. A compound as claimed in claim 1 which has the formula:



wherein X and Y are the same or different and are oxygen, sulfur, sulfone or sulfoxide; m and m' are the same or different and are integers in the range from 2 to 5; each R is  $\text{CH}_2\text{OH}$ ,  $\text{CHO}$  or  $\text{CO}_2\text{A}$  wherein A is a hydrogen atom, an alkyl radical having from 1 to 6 carbon atoms, an alkali or alkaline earth metal cation, an organic amine cation or an ammonium cation; each of  $\text{R}^1$  and  $\text{R}^2$ , which are the same or different, is an alkyl radical having from 1 to 6 carbon atoms, or  $\text{R}^1$  when taken together with the  $\text{R}^2$  which shares the same carbon atom represents  $-(\text{CH}_2)_{3-5}-$ ; and  $\text{R}^3$  is a hydrogen or halogen atom, an alkyl radical having from 1 to 6 carbon atoms, methoxy, phenyl, trifluoromethyl, 2,5-dichloro, 2,5-dimethyl, 2,3,5-trimethyl, carboxylic acyl or carboxylate alkyl of from 1 to 6 carbon atoms,

alpha-hydroxyalkyl of from 1 to 6 carbon atoms  
 2-chlorophenyl, 4-chlorophenyl, 2,4-dichlorophenyl,  
 4-methoxyphenyl, 2-tolyl, 4-tolyl or -D- wherein  
 D is O, CO, S, SO, SO<sub>2</sub>, CH<sub>2</sub>, or CHOH; provided that R<sup>3</sup> is  
 not a hydrogen atom when all the following conditions  
 are satisfied, namely when X and Y are both oxygen, m and m'  
 are both 2; R<sup>1</sup> and R<sup>2</sup> are both CH<sub>3</sub>, and R is CO<sub>2</sub>H;  
 or, when R is COOH, a pharmaceutically acceptable  
 salt of the compound.

3. A compound as claimed in claim 1, which has  
 the formula:



wherein L and M are the same or different and are oxygen,  
 sulfur, sulfone or sulfoxide; n and n' are the same or  
 different and are integers in the range from 2 to 5;  
 each of R<sup>4</sup> and R<sup>5</sup>, which are the same or different, is an  
 alkyl radical having from 1 to 6 carbon atoms; each R'  
 is CH<sub>2</sub>OH, CHO or CO<sub>2</sub>A wherein A is a hydrogen atom, an  
 alkyl radical having from 1 to 6 carbon atoms, an alkali or  
 alkaline earth metal cation, an organic amine cation or  
 an ammonium cation; R<sup>6</sup> is a hydrogen atom, 4-formyl,  
 2-nitro, 2-methyl, 5-methyl, 4-benzoyl or 4-chloro;  
 or, when R' is COOH, a pharmaceutically  
 acceptable salt of the compound.

4. A compound as claimed in claim 1 and which is  
 selected from: dimethyl 5,5'-[(2-fluoro-1,4-phenylene)bis  
 (oxy)]bis[2,2-dimethylpentanoate],  
 dimethyl 5,5'-[(2-acetyl-1,4-phenylene)bis(oxy)]-bis  
 [2,2-dimethylpentanoate],  
 dimethyl 5,5'-[(2-methyl-1,3-phenylene)bis(oxy)]-bis [2,2-  
 dimethylpentanoate],


-30-




5,5'-[(2'-chloro[1,1'-biphenyl]-2,5-diyl)bis(oxy)]bis-  
 [2,2-dimethylpentanoic acid],  
 5,5'-[(2'-methyl[1,1'-biphenyl]2,5-diyl)bis(oxy)]-  
 bis[2,2-dimethylpentanoic acid],  
 5,5'-[(2-trifluoromethyl-1,4-phenylene)bis(oxy)]-  
 bis[2,2-dimethylpentanoic acid],  
 dimethyl 5,5'-[[2-(1-oxopropyl)-1,4-phenylene]bis(oxy)]-  
 bis[2,2-dimethylpentanoate],  
 dimethyl 5,5'-[[2-(1-oxobutyl)-1,4-phenylene]bis(oxy)]-  
 bis[2,2-dimethylpentanoate],  
 dimethyl 5,5'-[(2-benzoyl-1,4-phenylene)bis(oxy)]-  
 bis[2,2-dimethylpentanoate],  
 dimethyl 5,5'[2,5-dimethyl-1,4-phenylenebis(oxy)]-bis[2,2-  
 dimethylpentanoate],  
 5,5'-[(2-phenylthio)-1,4-phenylene)bis(oxy)]bis[2,2-  
 dimethylpentanoic acid],  
 dimethyl 5,5'-[(2-methoxy-1,4-phenylene)bis(oxy)]-bis  
 [2,2-dimethylpentanoate],  
 dimethyl 5,5'-[(2-methyl-1,4-phenylene)bis(oxy)]-bis  
 [2,2-dimethylpentanoate],  
 dimethyl 5,5'[(2-bromo-1,4-phenylene)bis(oxy)]bis[2,2-  
 dimethylpentanoate],  
 5,5'-[(1,1'-biphenyl)-2,5-diylbis(oxy)]bis[2,2-  
 dimethylpentanoic acid],  
 5,5'-[1,4-phenylenebis(oxy)]bis[2,2-dimethyl pentanoic acid],  
 6,6'-[1,4-phenylenebis(oxy)]bis[2,2-dimethylhexanoic acid],  
 5,5'-[(2,5-dichloro-1,4-phenylene)bis(oxy)]bis-[2,2-  
 dimethylpentanoic acid],  
 4,4'-[(2-chloro-1,4-phenylene)bis(oxy)]bis-[2,2-  
 dimethylbutanoic acid],  
 5,5'-[(2-chloro-1,4-phenylene)bis(oxy)]bis-[2,2-  
 dimethylpentanoic acid],  
 5,5'-[1,3-phenylenebis(oxy)]bis[2,2-dimethylpentanoic acid],  
 4,4'-[1,3-phenylenebis(oxy)]bis[2,2-dimethylbutanoic acid],  
 6,6'-[1,3-phenylenebis(oxy)]bis[2,2-dimethylhexanoic acid],  
 7,7'-[1,2-phenylenebis(oxy)]bis[2,2-dimethylheptanoic acid],  
 5-[(4-[(4-carboxy-4-methylpentyl)oxy]phenyl)thio]-2,2-  
 dimethylpentanoic acid,




5- [[4-[(4-carboxy-4-methylpentyl)oxy]phenyl  
sulfinyl]-2,2-dimethylpentanoic acid,  
6,6'-[1,4-phenylenebis(oxy)] bis[2,2-dimethyl-1-hexanol],  
5,5'-[2-(hydroxyethyl)-1,4-phenylenebis(oxy)] bis-  
[2,2-dimethylpentanoic acid],  
5,5'-[[2-(benzyl)-1,4-phenylene]bis(oxy)] bis[2,2-  
dimethylpentanoic acid],  
5,5'-[2-(hydroxyphenylmethyl)-1,4-phenylenebis(oxy)]-  
bis[2,2-dimethylpentanoic acid],  
5,5'-[(2-hydroxymethyl)-1,4-phenylenebis(oxy)] bis[2,2-  
dimethylpentanoic acid],  
5,5'-[[2-(1-hydroxybutyl)-1,4-phenylene]bis(oxy)]-bis  
[2,2-dimethylpentanoic acid],  
5,5'-[[2(phenylsulfinyl)-1,4-phenylene]bis(oxy)][bis-  
[2,2-dimethylpentanoic acid], and  
5,5' [2-(1-hydroxypropyl)-1,4-phenylenebis(oxy)] bis-  
[2,2-dimethylpentanoic acid].

5. A process for preparing a compound as claimed  
in claim 1, which process comprises reacting a compound of  
the following general formula:-




wherein  $\textcircled{\text{Ar}}$  is , , or ; each of  
G and G' which are the same or different is oxygen  
or sulfur; and  $R_x$  represents any of the substituents  
 $R^3$  and  $R^6$  as defined in claim 1; with a compound of the  
following general formula:-




wherein Q is a leaving group and n, R,  $R^1$  and  $R^2$  are as  
defined in claim 1; and either isolating the resulting  
product or, when either of G and G' is sulfur, oxidizing  
the coupled product and isolating the oxidized product.

-32-

6. A process according to claim 5, which comprises reacting a compound of the formula




wherein R<sup>3</sup> is as defined in claim 1, with a compound of the formula




wherein Q is a leaving group and n, R, R<sup>1</sup> and R<sup>2</sup> are as defined in claim 1; and isolating the resulting product.

7. A process according to claim 5, which comprises reacting a compound of the formula




where R<sup>6</sup> is as defined in claim 1, with a compound of the formula







where Q is a leaving group and n, R, R<sup>1</sup> and R<sup>2</sup> are as defined in claim 1, and isolating the resulting product.

8. A process for producing a compound as claimed in claim 1, which process comprises reacting a compound of the following general formula:

-33-




wherein  $\text{Ar}$  is , , or ; each of  $G_1$  and  $G'_1$  which are the same or different is oxygen, sulfur, sulfone or sulfoxide;  $R_x$  represents any of the substituents  $R^3$  and  $R^6$  as defined in claim 1;  $Q$  and  $Q'$  are the same or different and are leaving groups; and  $m$  and  $m'$  are as defined in claim 1; with a compound of the following formula:




wherein  $R$ ,  $R^1$  and  $R^2$  are as defined in claim 1, and  $Z$  is an alkali metal; and isolating the resulting product.

9. A process according to claim 8, which comprises reacting a compound of the formula




wherein  $Q$  and  $Q'$  are leaving groups and  $m$ ,  $m'$  and  $R^6$  are as defined in claim 1; with a compound of the following formula




-34-

where  $R$ ,  $R^1$  and  $R^2$  are as defined in claim 1; and isolating the resulting product.

10. A process according to claim 8, which comprises reacting a compound of the formula




where  $Q$  and  $Q'$  are leaving groups and  $m$ ,  $m'$  and  $R^6$  are as defined in claim 1, with a compound of the following formula





where  $R$ ,  $R^1$  and  $R^2$  are as defined in claim 1, and isolating the resulting product.

CLAIMS (for Austria only)



1. A process for producing a compound having one of the following formulae 1 to 5:-




wherein X and Y are the same or different and are oxygen, sulfur, sulfone or sulfoxide; m and m' are the same or different and are integers in the range from 2 to 5; R is CH<sub>2</sub>OH, CHO or CO<sub>2</sub>A wherein A is a hydrogen atom, an alkyl radical having from 1 to 6 carbon atoms, an alkali or alkaline earth metal cation, an organic amine cation or an ammonium cation; each of R<sup>1</sup> and R<sup>2</sup> which are the same or different is an alkyl radical having from 1 to 6 carbon atoms, or R<sup>1</sup> when taken together with the R<sup>2</sup> sharing the same carbon atom represents -(CH<sub>2</sub>)<sub>3-5</sub>-; and R<sup>3</sup> is a hydrogen or halogen atom, an alkyl radical having from 1 to 6 carbon atoms, methoxy, phenyl, trifluoromethyl, 2,5-dichloro, 2,5-dimethyl, 2,3,5-trimethyl, carboxylic acyl or carboxylatoalkyl of from 1 to 6 carbon atoms, alpha-hydroxyalkyl having from 1 to 6 carbon atoms, 2-chlorophenyl, 4-chlorophenyl, 2,4-dichlorophenyl, 4-methoxyphenyl, 2-tolyl, 4-tolyl or -D- wherein D is O, CO, S, SO, SO<sub>2</sub>, CH<sub>2</sub> or CHOH; provided that R<sup>3</sup> is not a hydrogen atom when all of the following conditions are satisfied, namely when X and Y are both oxygen atoms, m and m' are both 2; R<sup>1</sup> and R<sup>2</sup> are both CH<sub>3</sub>, and R is CO<sub>2</sub>H:



wherein L and M are the same or different and are oxygen, sulfur, sulfone or sulfoxide; n and n' are the same or different and are integers in the range from 2 to 5; each of R<sup>4</sup> and R<sup>5</sup> which are the same or different is an alkyl radical having from 1 to 6 carbon atoms; each R' is CH<sub>2</sub>OH, CHO or CO<sub>2</sub>A wherein A is a hydrogen atom, an alkyl radical having from 1 to 6 carbon atoms, an alkali or alkaline earth metal cation, an organic amine cation or an ammonium cation; and R<sup>6</sup> is a hydrogen atom, 4-formyl, 2-nitro, 2-methyl, 5-methyl, 4-benzoyl or 4-chloro:







wherein, in formula 3, 4 or 5, each R'' is CH<sub>2</sub>OH, CHO or CO<sub>2</sub>A wherein A is a hydrogen atom, an alkyl radical having from 1 to 6 carbon atoms, an alkali or alkaline earth metal cation, an organic amine cation or an ammonium cation:

or, when R, R' or R'' is COOH in any such compound, a pharmaceutically acceptable salt of that compound:

which process comprises reacting a compound of the following general formula:




wherein (Ar) is , , or ;  
G and G' are the same or different and are oxygen or sulfur;  
and R<sub>x</sub> represents any of the substituents R<sup>3</sup> and R<sup>6</sup> as defined above; with a compound of the following general formula:-




wherein Q is a leaving group and n, R, R<sup>1</sup> and R<sup>2</sup> are as defined above; and either isolating the resulting product or, when either of G and G' is sulfur, oxidizing the coupled product and isolating the oxidized product.

2. A process according to claim 1, which comprises reacting a compound of the formula




where R<sup>3</sup> is as defined in claim 1, with a compound of the formula




wherein Q, n, R, R<sup>1</sup> and R<sup>2</sup> are as defined in claim 1; and isolating the resulting product.

3. A process according to claim 1, which comprises reacting a compound of the formula




where R<sup>6</sup> is as defined in claim 1, with a compound of the formula







where Q, n, R, R<sup>1</sup> and R<sup>2</sup> are as defined in claim 1, and isolating the resulting product.

4. A process for producing a compound as defined in claim 1, which process comprises reacting a compound of the following general formula:






wherein  $\text{Ar}$  is , , or ;  $G_1$  and  $G_1'$  are the same or different and are oxygen, sulfur, sulfone or sulfoxide;  $R_x$  represents any of the substituents  $R^3$  and  $R^6$  as defined in claim 1;  $Q$  and  $Q'$  are the same or different and are leaving groups; and  $m$  and  $m'$  are as defined in claim 1; with a compound of the following formula:-




wherein  $Z$  is an alkali metal and  $R$ ,  $R^1$  and  $R^2$  are as defined in claim 1; and isolating the resulting product.

5. A process according to claim 4, which process comprises reacting a compound of the following formula:




where  $Q$  and  $Q'$  are leaving groups and  $m$ ,  $m'$  and  $R^6$  are as defined in claim 1, with a compound of the formula'




where  $R$ ,  $R^1$  and  $R^2$  are as defined in claim 1; and isolating the resulting product.

6. A process according to claim 4, which process comprises reacting a compound of the formula



wherein  $Q$  and  $Q'$  are leaving groups and  $m$ ,  $m'$  and  $R^6$  are as defined in claim 1, with a compound of the formula



wherein  $R$ ,  $R^1$  and  $R^2$  are as defined in claim 1; and isolating the resulting product.

(12)

**EUROPEAN PATENT APPLICATION**

(21) Application number: 80304749.7

(22) Date of filing: 30.12.80

(51) Int. Cl.<sup>3</sup>: **A 61 K 31/225**

C 07 C 69/736, C 07 C 69/738  
C 07 C 149/40, C 07 C 79/46  
C 07 C 59/68, C 07 C 62/34  
C 07 C 147/11, C 07 C 43/205  
C 07 C 67/313, C 07 C 67/333

(30) Priority: 02.01.80 US 108978

(43) Date of publication of application:  
15.07.81 Bulletin 81/28

(88) Date of deferred publication of search report: 18.11.81

(84) Designated Contracting States:  
AT BE CH DE FR GB IT LI LU NL SE

(71) Applicant: **WARNER-LAMBERT COMPANY**  
201 Tabor Road  
Morris Plains, New Jersey 07950(US)

(72) Inventor: **Sircar, Ila**  
3615 Charter Place  
Ann Arbor Michigan 48105(US)






(74) Representative: **Jones, Michael Raymond et al.**  
Haseltine Lake & Co. 28 Southampton Buildings  
Chancery Lane  
London WC2A 1AT(GB)

(54) Novel substituted arylene compounds and processes for their production.

(57) Novel substituted arylene compounds and processes for their production are disclosed. These new compounds are useful as anti-arteriosclerotic agents. The novel compounds have the following formulae:-

**EP 0 032 063 A3**

./...



The radicals X, Y, L and M are each oxygen, sulfur, sulfone or sulfoxide and the other substituents are as defined at length in claim 1.



**THIS PAGE BLANK (USPTO)**

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**